

Listing all potential maximal cliques of a graph

Mochamad Suyudi, Sukono

Department of Mathematics, Faculty of Mathematics and Natural Sciences
Universitas Padjadjaran, Indonesia
moch.suyudi@gmail.com, sukono@unpad.ac.id

Mustafa Mamat

Faculty of Informatics and Computing, Universiti Sultan Zainal Abidin
Tembila Campus, 2200 Besut, Terengganu, Malaysia
musmat567@gmail.com

Abdul Talib Bon

Department of Production and Operations,
University Tun Hussein Onn Malaysia, Johor, Malaysia
talibon@gmail.com

Abstract

A potential maximal clique of a graph is a vertex set that induces a maximal clique in some minimal triangulation of that graph. It is known that if these objects can be listed in polynomial time for a class of graphs, the treewidth and the minimum fill-in are polynomially tractable for these graphs. We show here that the potential maximal cliques of a graph can be generated in polynomial time in the number of minimal separators of the graph. Thus, the treewidth and the minimum fill-in are polynomially tractable for all classes of graphs with a polynomial number of minimal separators.

Keywords

Graph algorithms, Treewidth, Minimal separators, Potential maximal cliques.

1. Introduction

The notion of treewidth was introduced at the beginning of the 1980s by (Robertson and Seymour 1984), (Robertson and Seymour 1986), in the framework of their graph minor theory. A graph H is a minor of a graph G if we can obtain H from G by repeatedly using the following operations: vertex deletion, edge deletion and edge contraction. Among the deep results obtained by Robertson and Seymour, we can cite the fact that every class of graphs closed under taking minors which does not contain all the planar graphs has bounded treewidth.

A graph is chordal or triangulated if every cycle of length greater or equal to four has a chord, i.e. an edge between two non-consecutive vertices of the cycle. A triangulation of a graph is a chordal embedding, that is a supergraph, on the same vertex set, which is triangulated. The treewidth problem consists in finding a triangulation such that the maximum size of a clique is as small as possible. Another close problem is the minimum fill-in problem. Here we have to find a triangulation of the graph such that the number of added edges is minimum. In both cases, we can restrict to minimal triangulations, i.e. triangulations with a set of edges minimal by inclusion.

The treewidth and the minimum fill-in play an important role in various areas of computer science. Among the real-life applications we can cite sparse matrix factorization (Rose 1970), Cholesky factorization (Liu 1996), perfect phylogeny problem (Kannan and Warnow 1990) and register allocation (Thorup 1998), (Bodlaender et al. 1998). General techniques exist for solving classical algorithmic graph theory problems for graphs of bounded treewidth (Arnborg and Proskurowski 1989), (Courcelle and Mosbah 1993), (Arnborg et al. 1993), (Bodlaender and de Fluiter 1996). For an extensive survey of these applications see also (Bodlaender 1993), (Bodlaender 1997).

For computing the treewidth we usually look for a tree decomposition, that is a tree such that each node of the tree is labeled by a vertex set of the graph. The labels of the nodes must respect some constraints: every vertex of the graph must appear in some label, the endpoints of an edge must appear in a same label, if a same

vertex is in two different labels it must be in all the labels on the unique path of the tree connecting the two occurrences of the vertex. The width of the tree decomposition is then the size of the largest label minus one, and the treewidth is the smallest width over all the tree decompositions of the graph. In some sense, the smaller the treewidth of a graph, the more it has tree-like structure. Many graph problems that model real-life problems are intractable in the sense that they are NP-hard. If we deal with a class of graphs with bounded treewidth most of these problems become polynomial and even linear e.g. maximum independent set, hamiltonian circuit or Steiner tree. There are two ways to solve problems when the treewidth is bounded, the first uses dynamic programming (Bodlaender 1993), (Hagerup 1997) and the second is based upon reduction techniques (Arnborg et al. 1993), (Bodlaender and de Fluiter 1996).

Unfortunately the computation of the treewidth and of the minimum fill-in of a graph are NP-hard (Arnborg et al. 1987), (Yannakakis 1981) even for co-bipartite graphs. However, a polynomial time approximation algorithm with $O(\log n)$ performance ratio is described in (Bodlaender et al. 1995). The problem of the existence of a polynomial algorithm to approximate the treewidth of a graph within a multiplicative constant remains still open. For any fixed constant k , there exist polynomial algorithms finding a tree decomposition of width at most k if such a decomposition exists. (Arnborg et al. 1987) gave the first algorithm that solves this problem in $O(n^{k+2})$ time. Since numerous improvements have been done on the domain until the linear time algorithm of (Bodlaender 1996). Notice that the constant hidden by the O notation is doubly exponential in k^2 . Several results for treewidth appeared in the literature in connection with logic. The works by (Arnborg et al. 1993), (Courcelle 1992), (Courcelle and Mosbah 1993) led to the conclusion that all the problems which are expressible in extended monadic second order logic can be solved in linear time for graphs of bounded treewidth.

There exist several classes of graphs with unbounded treewidth for which we can solve polynomially the problem of the treewidth and the minimum fill-in. Among them there are the chordal bipartite graphs (Kloks and Kratsch 1995), (Chang 1996), circle and circular-arc graphs (Sundaram et al. 1994), (Kloks et al. 1998), AT-free graphs with polynomial number of *minimal separators* (Kloks et al. 1997). Most of these algorithms use the fact that these classes of graphs have a polynomial number of minimal separators, i.e. for each of these classes, there is a polynomial $P(n)$ such that every graph G in the class of n vertices has at most $P(n)$ minimal separators. It was conjectured in (Kloks et al. 1993), (Kloks et al. 1994) that the treewidth and the minimum fill-in should be tractable in polynomial time for every graph class having a polynomial number of minimal separators. We solve here this ESA'93 conjecture.

The crucial interplay between the minimal separators of a graph and the minimal triangulations was pointed out by (Kloks et al. 1995), these results were concluded in (Parra and ScheLer 1997). Two minimal separators S and T cross if T intersects two connected components of $G \setminus S$, otherwise they are parallel. The result of (Parra and ScheLer 1997) states that a minimal triangulation is obtained by considering a maximal set of pairwise parallel separators and by completing them i.e. by adding all the missing edges inside each separator. However this characterization gives no algorithmic information about how we should construct a minimal triangulation in order to minimize the clique size or the fill-in.

Trying to solve this later conjecture, we studied in (Bouchitté and Todinca 1998), (Bouchitté and Todinca 1999) the notion of a potential maximal clique. A vertex set K is a potential maximal clique if it appears as a maximal clique in some minimal triangulation. In (Bouchitté and Todinca 1998), we characterized a potential maximal clique in terms of the maximal sets of neighbor separators, which are the minimal separators contained in it. We designed an algorithm which takes as input the graph and the maximal sets of neighbor separators and which computes the treewidth in polynomial time in the size of the input. For all the classes mentioned above we can list the maximal sets of neighbor separators in polynomial time, so we unified all the previous algorithms. Actually, the previous algorithms compute the maximal sets of neighbor separators in an implicit manner. In (Bouchitté and Todinca 1999), we gave a new characterization of the potential maximal cliques avoiding the minimal separators. This allowed us to design a new algorithm that, given a graph and its potential maximal cliques, computes the treewidth and the minimum fill-in in polynomial time. Moreover this approach permitted us to solve the two problems for a new class of graphs, namely the weakly triangulated graphs. It was probably the last natural class of graphs with polynomial number of minimal separators for which the two problems remained open.

This paper is devoted to solve the ESA'93 conjecture, that is the treewidth and the minimum fill-in are polynomially tractable for every graph class having a polynomial number of minimal separators. Recall that if we are able to generate all the potential maximal cliques of any graph in polynomial time in the number of its minimal separators, then the treewidth and the minimum fill-in are also computable in polynomial time in the number of minimal separators. We define the notion of an active separator for a potential maximal clique which leads to two results. First, the number of potential maximal cliques is polynomially bounded by the number of minimal separators. Secondly, we are able to enumerate the potential maximal cliques in polynomial time in their number. These results reinforce our conviction that the potential maximal cliques are the pertinent objects to study when dealing with treewidth and minimum fill-in.

2. Preliminaries

Throughout this paper, we consider finite, simple, undirected and connected graphs.

Let $G = (V, E)$ be a graph. We will denote by n and m the number of vertices, respectively the number of edges of G . For a vertex set $V' \subseteq V$ of G , we denote by $N_G(V')$ the neighborhood of V' in $G[V']$ – so $N_G(V') \subseteq V \setminus V'$.

A subset $S \subseteq V$ is an a, b -separator for two nonadjacent vertices $a, b \in V$ if the removal of S from the graph separates a and b in different connected components. S is a minimal a, b -separator if no proper subset of S separates a and b . We say that S is a minimal separator of G if there are two vertices a and b such that S is a minimal a, b -separator. Notice that a minimal separator can be strictly included in another one. We denote by Δ_G the set of all minimal separators of G .

Let G be a graph and S a minimal separator of G . We note $\mathcal{C}_G(S)$ the set of connected components of $G \setminus S$. A component $C \in \mathcal{C}_G(S)$ is a full component associated to S if every vertex of S is adjacent to some vertex of C , i.e. $N_G(C) = S$. The following lemmas see (Golombic 1980) for a proof) provide different characterizations of a minimal separator.

Lemma 1. A set S of vertices of G is a minimal a, b -separator if and only if a and b are in different full components of S .

Lemma 2. Let G be a graph and S be an a, b -separator of G . Then S is a minimal a, b -separator if and only if for any vertex x of S there is a path from a to b that intersects S only in x .

If $C \in \mathcal{C}_G(S)$, we say that $(S, C) = S \cup C$ is a block associated to S . A block (S, C) is called full if C is a full component associated to S .

Let now $G = (V, E)$ be a graph and $G' = G[V']$ an induced subgraph of G . We will compare the minimal separators of G and G' .

Lemma 3. Let G be a graph and $V' \subset V$ a vertex set of G . If S is a minimal a, b -separator of the induced subgraph $G' = G[V']$; then there is a minimal a, b -separator T of G such that $T \cap V' = S$.

Proof. Let $S' = S \cup (V \setminus V')$. Clearly, S' is an a, b -separator in G . Let T be any minimal a, b separator contained in S' . We have to prove that $S \subseteq T$. Let x be any vertex of S and suppose that $x \notin T$. Since S is a minimal a, b -separator of G' , we have a path μ joining a and b in G' that intersects S only in x (see Lemma 2). But μ is also a path of G , that avoids T , contradicting the fact that T is an a, b -separator. It follows that $S \subseteq T$. Clearly, $T \cap V' \subseteq S$ by construction of T , so $T \cap V' = S$.

The next corollary follows directly from Lemma 3.

Corollary 4. Let $G = (V, E)$ be a graph and a be a vertex of G . Consider the graph $G' = G[V \setminus \{a\}]$. Then for any minimal separator S of G' , we have that S or $S \cup \{a\}$ is a minimal separator of G . In particular, $|\Delta_G| \geq |\Delta_{G'}|$.

3. Potential maximal cliques and maximal sets of neighbor separators

The potential maximal cliques are the central object of this paper. We present in this section some known results about the potential maximal cliques of a graph (see also (Bouchitté and Todinca 1998), (Bouchitté and Todinca 1999), (Todinca 1999)).

Definition 5. A vertex set Ω of a graph G is called a potential maximal clique if there is a minimal triangulation H of G such that Ω is a maximal clique of H .

We denote by Π_G the set of potential maximal cliques of the graph G .

A potential maximal clique Ω is strongly related to the minimal separators contained in Ω . In particular, any minimal separator of G is contained in some potential maximal clique of G and a potential maximal clique contains at most n minimal separators, where n is the size of G (by Theorem 8). of potential maximal cliques of G is at least $|\Delta_G|/n$. If K is a vertex set of G , we denote by $\Delta_G(K)$ the minimal separators of G included in K .

Definition 6. A set \mathcal{S} of minimal separators of a graph G is called maximal set of neighbor separators if there is a potential maximal clique Ω of G such that $\mathcal{S} = \Delta_G(\Omega)$. We also say that \mathcal{S} borders Ω in G .

We proved in (Bouchitté and Todinca 1999) that the potential maximal cliques of a graph are sufficient for computing the treewidth and the minimum fill-in of that graph.

Theorem 7 (Bouchitté and Todinca 1999). *Given a graph G and its potential maximal cliques Π_G , we can compute the treewidth and the minimum fill-in of G in $O(n^2|\Delta_G| \times |\Pi_G|)$ time.*

Let now K be a set of vertices of a graph G . The connected components of $G \setminus K$ will be denoted by $C_1(K), \dots, C_p(K)$. We denote by $S_i(K)$ the vertices of K adjacent to at least one vertex of $C_i(K)$. When no confusion is possible we will simply speak of C_i and S_i . If $S_i(K) = K$ we say that $C_i(K)$ is a *full component* associated to K . Finally, we denote by $\mathcal{E}_G(K)$ the set of all $S_i(K)$ in the graph G , i.e. $\mathcal{E}_G(K)$ is formed by the neighborhoods, in the graph G , of the connected components of $G \setminus K$.

Consider graph $G = (V, E)$ and a vertex set $X \subseteq V$. We denote by G_X the graph obtained from G by completing X , i.e. by adding an edge between every pair of nonadjacent vertices of X . If $\mathcal{X} = \{X_1, \dots, X_p\}$ is a set of subsets of V , $G_{\mathcal{X}}$ is the graph obtained by completing all the elements of \mathcal{X} .

Theorem 8 (Bouchitté and Todinca 1999). *Let $K \subseteq V$ be a set of vertices. K is a potential maximal clique if and only if:*

- (1) $G \setminus K$ has no full components associated to K .
- (2) $G_{\mathcal{E}_G(K)}[K]$ is a clique.




Fig. 1. Potential maximal cliques.

Moreover, if K is a potential maximal clique, then $\mathcal{E}_G(K)$ is the maximal set of neighbor separators bordering K ; i.e. $\mathcal{E}_G(K) = \Delta_G(K)$.

For example, in Fig. 1, the vertex sets $\{b, c, e, g\}$ and $\{b, d, e\}$ are potential maximal cliques of the graph of Fig. 1a and the vertices $\{x, y, z, t\}$ form a potential maximal clique of the graph of Fig. 1b.

Remark 9. If K is a potential maximal clique of G , for any pair of vertices x and y of K either x and y are adjacent in G or they are connected by a path entirely contained in some C_i of $G \setminus K$ except for x and y . The second case comes from the fact that if x and y are not adjacent in G they must belong to the same S_i to ensure that K becomes a clique after the completion of $\mathcal{E}_G(K)$. When we will refer to this property we will say that x and y are connected via the connected component C_i .

Remark 10. Consider a minimal separator S contained in a potential maximal clique Ω . Let us compare the connected components of $G \setminus S$ and the connected components of $G \setminus \Omega$ (see (Bouchitté and Todinca 1999) for the proofs). The set $\Omega \setminus S$ is contained in a full component C_Ω associated to S . All the other connected components of $G \setminus S$ are also connected components of $G \setminus \Omega$. Conversely, a connected component C of $G \setminus \Omega$ is either a connected component of $G \setminus S$ (in which case $N_G(C) \subseteq S$) or it is contained in C_Ω (in which case $N_G(C) \not\subseteq S$).

Remark 11. Unlike the minimal separators, a potential maximal clique Ω' cannot be strictly included in another potential maximal clique Ω . Indeed, for any proper subset Ω' of a potential maximal clique Ω , the difference $\Omega \setminus \Omega'$ is in a full component associated to Ω' .

Theorem 8 leads to a polynomial algorithm that, given a vertex set of a graph G , decides if K is a potential maximal clique of G .

Corollary 12. Given a vertex set K of a graph G ; we can recognize in $O(nm)$ time if K is a potential maximal clique of G .

Proof. We can compute in linear time the connected components C_i of $G \setminus K$ and their neighborhoods S_i . We can also verify in linear time that $G \setminus K$ has no full components associated to K .

For each $x \in K$, we compute all the vertices $y \in K$ that are adjacent to x in G or connected to x via a C_i in linear time (we have to search the neighborhood of x and the connected components C_i with $x \in S_i$). So we can verify in $O(nm)$ time if K satisfies the conditions of Theorem 8.

4. Potential maximal cliques and active separators

Theorem 8 tells us that if Ω is a potential maximal clique of a graph G , then Ω is a clique in $G_{\Delta_G(\Omega)}$. We will divide the minimal separators of $\Delta_G(\Omega)$ into two classes: those which create edges in $G_{\Delta_G(\Omega)}$, which are called *actives*, and the others, which are called *inactives*. More precisely:

Definition 13. Let Ω be a potential maximal clique of a graph G and let $S \subset \Omega$ be a minimal separator of G . We say that S is an active separator for Ω if Ω is not a clique in the graph $G_{\Delta_G(\Omega) \setminus \{S\}}$, obtained from G by completing all the minimal separators contained in Ω , except S . Otherwise, S is called inactive for Ω .

Proposition 14. Let Ω be a potential maximal clique of G and $S \subset \Omega$ a minimal separator; active for Ω . Let (S, C_Ω) be the block associated to S containing Ω and let $x, y \in \Omega$ be two non-adjacent vertices of $G_{\Delta_G(\Omega) \setminus \{S\}}$. Then $\Omega \setminus S$ is an minimal x, y -separator in $G[C_\Omega \cup \{x, y\}]$.

Proof. Remark that the vertices x and y , non-adjacent in $G_{\Delta_G(\Omega) \setminus \{S\}}$, exist by definition of an active separator. Moreover, since $G_{\Delta_G(\Omega)}$ is a clique, we must have $x, y \in S$.

Let us prove first that $\Omega \setminus S$ is a x, y -separator in the graph $G' = G[C_\Omega \cup \{x, y\}]$. Suppose that x and y are in a same connected component C_{xy} of $G' \setminus (\Omega \setminus S)$. Let $C = C_{xy} \setminus \{x, y\}$. Clearly, $C \subset C_\Omega$ is a connected component of $G \setminus \Omega$. Let T be the neighborhood of C in G . By Theorem 8, T is a minimal separator of G , contained in Ω . By construction of T , we have $x, y \in T$. Notice that $T \neq S$, otherwise S would separate C and Ω , contradicting the fact that $C \subset C_\Omega$ (see Remark 10). It follows that T is a minimal separator of $\Delta_G(\Omega)$, different from S and containing x and y . This contradicts the fact that x and y are not adjacent in $G_{\Delta_G(\Omega) \setminus \{S\}}$. We can conclude that $\Omega \setminus S$ is an x, y -separator of G' .

We prove now that $\Omega \setminus S$ is a minimal x, y -separator of G' . We will show that, for any vertex $z \in \Omega \setminus S$, there is a path μ joining x and y in G' and such that μ intersects $\Omega \setminus S$ only in z . By Theorem 8, x and z are adjacent in $G_{\Delta_G(\Omega)}$, so x and z are adjacent in G or they are connected via a connected component C_i of $G \setminus \Omega$. Notice that $C_i \subset C_\Omega$: indeed, if $C_i \not\subset C_\Omega$, then C_i will be contained in some connected component D of $G \setminus S$, different from C_Ω . According to Remark 10, we would have $N_G(C_i) \subseteq N_G(D) \subseteq S$, contradicting $z \in S_i$. In both cases we have a path μ' from x to z in G' , that intersects $\Omega \setminus S$ only in z .

For the same reasons, z and y are adjacent in G , or there is a connected component C_j of $G \setminus \Omega$ such that $C_j \subset C_\Omega$ and $z, y \in S_j = N_G(C_j)$. This gives us a path μ'' from z to y in G' , such that $\mu'' \cap (\Omega \setminus S) = \{z\}$. Remark that $C_i \neq C_j$, otherwise we would have a path from x to y in $C_i \cup \{x, y\}$, contradicting the fact that $\Omega \setminus S$ separates x and y in G' . So the paths μ' and μ'' are disjoint except for z , and their concatenation is a path μ , joining x and y in G' and intersecting $\Omega \setminus S$ only in z . We conclude by Lemma 2 that $\Omega \setminus S$ is a minimal separator of G' .

By Proposition 14, the set $T' = \Omega \setminus S$ is a minimal separator of the subgraph of G induced by $C_\Omega \cup \{x, y\}$. By Lemma 3, there is a separator T of G such that $T' \subseteq T$ and $T \cap C_\Omega = T'$. We deduce:

Theorem 15. Let Ω be a potential maximal clique and S be a minimal separator; active for Ω . Let (S, C_Ω) be the block associated to S containing Ω . There is a minimal separator T of G such that $\Omega = S \cup (T \cap C_\Omega)$.

It follows easily that the number of potential maximal cliques containing at least one active separator is polynomially bounded in the number of minimal separators of G . More exactly the number of these potential maximal cliques is bounded by the number of blocks (S, C_Ω) multiplied by the number of minimal separators T , so by $n|\Delta_G|^2$. Clearly, these potential maximal cliques have a simple structure and can be computed directly from the minimal separators of the graph.




Fig. 2. Active and inactive separators.

Nevertheless, a potential maximal clique may not have active separators. For example in Fig. 2, the potential maximal clique $\Omega = \{a, c, f, h\}$ contains the minimal separators $\{a, f, h\}$, $\{c, f, h\}$, $\{a, c, f\}$ and $\{a, c, h\}$, but no one of them is active for Ω . Let us make a first observation about the potential maximal cliques containing inactive minimal separators.

Proposition 16. *Let Ω be a potential maximal clique and $S \subset \Omega$ a minimal separator which is inactive for Ω . Let D_1, \dots, D_p be the full components associated to S that do not intersect Ω . Then Ω is a potential maximal clique of the graph $G \setminus \bigcup_{i=1}^p D_i$.*

Proof. Let $G' = G \setminus \bigcup_{i=1}^p D_i$. The connected components of $G' \setminus \Omega$ are exactly the connected components of $G \setminus \Omega$, except for D_1, \dots, D_p , and their neighborhoods in G' are the same as in G . It follows that the set $\mathcal{E}_{G'}(\Omega)$ of the neighborhoods of the connected components of $G' \setminus \Omega$ is exactly $\Delta_G(\Omega) \setminus \{S\}$. Clearly, $G' \setminus \Omega$ has no full components associated to Ω . Since S is not active for Ω , we deduce that Ω is a clique in $G' \setminus \mathcal{E}_{G'}(\Omega)$. So, by Theorem 8, Ω is a potential maximal clique of G' .

5. Removing a vertex

Let $G = (V, E)$ be a graph and a be a vertex of G . We denote by G' the graph obtained from G by removing a , i.e. $G' = G[V \setminus \{a\}]$. We will show here how to obtain the potential maximal cliques of G using the minimal separators of G and G' and the potential maximal cliques of G' . By Corollary 4, we know that G has at least as many minimal separators as G' : for any minimal separator S of G' , either S is a minimal separator of G , or $S \cup \{a\}$ is a minimal separator of G . It will follow that the potential maximal cliques of a graph can be computed in polynomial time in the size of the graph and the number of its minimal separators.

Proposition 17. *Let Ω be a potential maximal clique of G such that $a \in \Omega$. Then $\Omega = \Omega' \setminus \{a\}$ is either a potential maximal clique of G' or a minimal separator of G .*

Proof. Let C_1, \dots, C_p be the connected components of $G \setminus \Omega$ and S_i be the neighborhood of C_i in G . We denote as usual by $\mathcal{E}_G(\Omega)$ the set of all the S_i 's. Remark that the connected components of $G' \setminus (\Omega \setminus \{a\})$ are exactly C_1, \dots, C_p and their neighborhoods in G' are respectively $S_1 \setminus \{a\}, \dots, S_p \setminus \{a\}$. Since Ω is a clique in $G \setminus \mathcal{E}_G(\Omega)$ (by Theorem 8), it follows that $\Omega = \Omega' \setminus \{a\}$ is a clique in $G' \setminus \mathcal{E}_{G'}(\Omega)$. If $G' \setminus \Omega'$ has no full components associated to Ω' , then Ω' is a potential maximal clique of G' , according to Theorem 8. Suppose now that C_1 is a full component associated to Ω in G' . Since C_1 is not a full component associated to Ω in G , it follows that $N_G(C_1) = \Omega'$. Thus, Ω' is a minimal separator of G , by Theorem 8.

Lemma 18. *Let G be a graph and \tilde{G} be any induced subgraph of G . Consider a potential maximal clique Ω of \tilde{G} . Suppose that for any connected component C of $G \setminus \tilde{G}$, its neighborhood $N_G(C)$ is strictly contained in Ω . Then Ω is also a potential maximal clique of G .*

Proposition 19. *Let Ω be a potential maximal clique of G such that $a \notin \Omega$. Let C_a be the connected component of $G \setminus \Omega$ containing a and let S be the minimal separator of Ω such that $S = N(C_a)$. If Ω is not a potential maximal clique of $G' = G[V \setminus \{a\}]$, then S is active for Ω . Moreover, S is not a minimal separator of G' .*

Theorem 20. *Let Ω be a potential maximal clique of G and let $G' = G[G \setminus \{a\}]$. Then one of the following cases holds:*

- (1) $\Omega = \Omega' \cup \{a\}$, where Ω' is a potential maximal clique of G' .
- (2) $\Omega = \Omega'$, where Ω' is a potential maximal clique of G' .
- (3) $\Omega = S \cup \{a\}$, where S is a minimal separator of G .
- (4) $\Omega = S \cup (C \cap T)$, where S is a minimal separator of G , C is a connected component of $G \setminus S$ and T is a minimal separator of G . Moreover; S does not contain a and S is not a minimal separator of G' .

Corollary 21. Let G be a graph, a be a vertex of G and $G' = G \setminus \{a\}$. The number $|\Pi_G|$ of potential maximal cliques of G is polynomially bounded in the number $|\Pi_{G'}|$ of potential maximal cliques of G' ; the number $|\Delta_G|$ of minimal separators of G and the size n of G .

More precisely; $|\Pi_G| \leq |\Pi_{G'}| + n(|\Delta_G| - |\Delta_{G'}|)|\Delta_G| + |\Delta_G|$.

Let now a_1, a_2, \dots, a_p be an arbitrary ordering of the vertices of G . We denote by G_i the graph $G[\{a_1, \dots, a_i\}]$, so $G_n = G$ and G_1 has a single vertex. By Corollary 21 we have that for any $i, 1 \leq i \leq n$, $|\Pi_{G_{i+1}}| \leq |\Pi_{G_i}| + n(|\Delta_{G_{i+1}}| - |\Delta_{G_i}|)|\Delta_{G_{i+1}}| + |\Delta_{G_{i+1}}|$. Notice that $|\Delta_{G_i}| \leq |\Delta_{G_{i+1}}|$, in particular each graph G_i has at most $|\Delta_G|$ minimal separators.

Clearly, the graph G_1 has a unique potential maximal clique. It follows directly that the graph G has at most $n|\Delta_G|^2 + n|\Delta_G| + 1$ potential maximal cliques.

Proposition 22. The number of the potential maximal cliques of a graph is polynomially bounded in the number of its minimal separators and in the size of the graph.

More precisely, a graph G has at most $n|\Delta_G|^2 + n|\Delta_G| + 1$ potential maximal cliques.

We give now an algorithm computing the potential maximal cliques of a graph. We suppose that we have a function $IS_PMC(\Omega, G)$, that returns *TRUE* if Ω is a potential maximal clique of G , *FALSE* otherwise.

The function ONE_MORE_VERTEX computes the potential maximal cliques of a graph G from the potential maximal cliques of a graph $G' = G \setminus \{a\}$. This function is based on Theorem 20. The main program successively computes the potential maximal cliques of the graphs $G_i = G[\{a_1, \dots, a_i\}]$. Notice that we can compute the vertex ordering such that each of the graphs G_i is connected.

Theorem 23. The potential maximal cliques of a graph can be listed in polynomial time in its size and the number of its minimal separators.

More exactly; the potential maximal cliques of a graph are computable in $O(n^2 m |\Delta_G|^2)$ time.

Proof. Let us analyze the complexity of the algorithm. The sets of vertex sets, like Π_G and Δ_G , will be represented by trees, in such manner that the adjunction of a new element and testing that a vertex set belongs to our set will be done in linear time (see for example (Kloks et al. 1998)). We also know by Corollary 12 that a call of the function IS_PMC takes $O(nm)$ time.

Let us start with the function ONE_MORE_VERTEX . We compute the cost of one execution of this function.

The cost of the first for loop is at most $|\Pi_{G'}|nm$. But we can strongly reduce this complexity, using a different test for verifying that Ω' , respectively $\Omega' \cup \{a\}$ are potential maximal cliques of G . Suppose that we want to check if a potential maximal clique Ω' of G' is also a potential maximal clique of G . Any connected component C' of Ω' is contained in some connected component C of $G \setminus \Omega$ and we have $N_{G'}(C') \subseteq N_G(C)$. Since Ω' is a clique in the graph $G_{\setminus G'(\Omega')}$, Ω' is a clique in the graph $G_{\setminus G(\Omega')}$. Therefore, all we have to check is that $G \setminus \Omega'$ has no full connected components associated to Ω' , which can be done in linear time. Suppose now that Ω' is a potential maximal clique of G' and let us verify if $\Omega = \Omega' \cup \{a\}$ is a potential maximal clique of G . Clearly, the connected components of $G \setminus \Omega$ are the same as the connected components of $G' \setminus \Omega'$. The neighborhood $N_G(C)$ of a connected component

function ONE_MORE_VERTEX

Input: the graphs G, G' and a vertex a such that $G' = G \setminus \{a\}$;
the potential maximal cliques $\Pi_{G'}$ of G' , the minimal separators
 $\Delta_{G'}, \Delta_G$ of G' and G .

Output: the potential maximal cliques Π_G of G .

begin

$\Pi_G \leftarrow \emptyset$

```

for each p.m.c.  $\Omega' \in \Pi_{G'}$ 
  if  $IS\_PMC(\Omega', G)$  then
     $\Pi_G \leftarrow \Pi_G \cup \{\Omega'\}$ 
  else
    if  $IS\_PMC(\Omega' \cup \{a\}, G)$  then
       $\Pi_G \leftarrow \Pi_G \cup \{\Omega' \cup \{a\}\}$ 
for each minimal separator  $S \in \Delta_G$ 
  if  $IS\_PMC(S \cup \{a\}, G)$  then
     $\Pi_G \leftarrow \Pi_G \cup \{S \cup \{a\}\}$ 
  if ( $a \notin S$  and  $S \notin \Delta_{G'}$ ) then
    for each  $T \in \Delta_G$ 
      for each full component  $C$  associated to  $S$  in  $G$ 
        if  $IS\_PMC(S \cup (T \cap C), G)$  then
           $\Pi_G \leftarrow \Pi_G \cup \{S \cup (T \cap C)\}$ 
return  $\Pi_G$ 
end

```

of $G \setminus \Omega$ is either $N_{G'}(C)$ or $N_{G'}(C) \cup \{a\}$. It follows that $G \setminus \Omega'$ has no full components associated to Ω and that any two vertices $x, y \in \Omega'$ are adjacent in $G_{\mathcal{G}(G, \Omega)}$. It remains to check that, in the graph $G_{\mathcal{G}(G, \Omega)}$, a is adjacent to any vertex $x \in \Omega'$. This test can be done in linear time: by searching $N_G(a)$ and the connected components C_i of $G \setminus \Omega$ with $a \in S_i$, we compute the vertices of Ω' adjacent to a in G or connected to a via C_i . We conclude that the cost of the first for loop is $O(m|\Pi_{G'}|)$, where $|\Pi_{G'}| = O(n|\Delta_G|^2)$.

In the second for loop, computing the potential maximal cliques of type 3, i.e. of type $S \cup \{a\}$, costs $O(nm|\Delta_G|)$ time. This is due to the cost of the Δ_G calls to function IS_PMC . Remark that here we could also test in linear time if $\Omega = S \cup \{a\}$ is a potential maximal clique of G . Since $S \subseteq N_G(C)$ for some

main program

Input: a graph G

Output: the potential maximal cliques Π_G of G

begin

let $\{a_1, \dots, a_n\}$ be the vertices of G

$\Pi_{G_1} \leftarrow \{\{a_1\}\}$

$\Delta_{G_1} \leftarrow \emptyset$

for $i = 1, n-1$

compute $\Delta_{G_{i+1}}$

$\Pi_{G_{i+1}} = ONE_MORE_VERTEX(G_i, G_{i+1}, \Pi_{G_i}, \Delta_{G_i}, \Delta_{G_{i+1}})$

$\Pi_G = \Pi_{G_n}$

end

connected component of $G \setminus \Omega$ (see proof of Proposition 17), we only have to test that $G \setminus \Omega$ has no full components associated to Ω and that a is adjacent in $G_{\mathcal{G}(G, \Omega)}$ to every $x \in S$. Anyway, this will not change the global complexity of the algorithm.

The call to function IS_PMC in the inner loop is done $n|\Delta_G|(|\Delta_G| - |\Delta_{G'}|)$ times. Indeed, we have shown in the proof of Corollary 21 that the number of minimal separators $S \in \Delta_G$ such that $a \notin S$ and $S \notin \Delta_{G'}$ is at most $|\Delta_G| - |\Delta_{G'}|$. The number of iteration of the second and third loop are clearly $|\Delta_G|$ and respectively n . So the cost of all the calls to function IS_PMC will be $O(n^2m|\Delta_G|(|\Delta_G| - |\Delta_{G'}|))$. So one execution of the function ONE_MORE_VERTEX takes at most $O(nm|\Delta_G|^2 + n^2m|\Delta_G|(|\Delta_G| - |\Delta_{G'}|))$ time.

We can compute now the complexity of the main program. Computing the minimal separators of a graph G can be done in $O(n^3|\Delta_G|)$ time, using the algorithm of Berry et al. [4]. If we do this calculus one time for each graph G_i , this would take $O(n^4|\Delta_G|)$. But notice that each graph G_i is an induced subgraph of G . Consequently, for each minimal separator S_i of G_i , there is a minimal separator S of G such that $S_i = S \cap \{a_1, \dots, a_i\}$. We can compute first the minimal separators of the input graph G , in $O(n^3|\Delta_G|)$ time. For computing the minimal separators of a graph G_i , we will take each $S \in \Delta_G$ and we will verify if $S_i = S \cap \{a_1, \dots, a_i\}$ is a minimal separator of G_i . A verification of type $S_i \in \Delta_{G_i}$ can be done in linear time: it is sufficient to test that $G_i \setminus S_i$ has at least two full components associated to S_i (see Lemma 1). Therefore, computing the minimal separators of all the graphs G_i will not exceed $O(n^3|\Delta_G|)$ steps.

Remember that the i th call of the function *ONE_MORE_VERTEX* costs at most $O(nm|\Delta_{G_{i+1}}|^2 + n^2m|\Delta_{G_{i+1}}|(|\Delta_{G_{i+1}}| - |\Delta_{G_i}|))$ time. Using the fact that for all i we have that $|\Delta_{G_i}| \leq |\Delta_G|$, it follows that the n calls of the function *ONE_MORE_VERTEX* will take $O(n^2m|\Delta_G|^2)$ steps.

We conclude that the global complexity of the algorithm is $O(n^2m|\Delta_G|^2)$.

We deduce directly from Theorem 7, Proposition 22 and Theorem 23:

Theorem 24. *The treewidth and the minimum fill-in of a graph can be computed in polynomial time in the size of the graph and the number of its minimal separators. The complexity of the algorithm is $O(n^3|\Delta_G|^3 + n^2m|\Delta_G|^2)$.*

6. Conclusion

The notion of a potential maximal clique seems to be very useful for the study of the treewidth and the minimum fill-in problems. We proved in (Bouchitté and Todinca 1999) that the potential maximal cliques are sufficient for computing the treewidth and the minimum fill-in of a graph. In this paper, we enumerate the potential maximal cliques in polynomial time in the number of minimum separators of the input graph. In particular, this gives a polynomial algorithm computing the treewidth and the minimum fill-in for all the classes of graphs with polynomial number of minimal separators.

Our generic algorithm is, of course, less efficient than algorithms working on particular graph classes. While enumerating the potential maximal cliques of the input graph, we also generate sets of vertices that do not form potential maximal cliques, so we lose time. There might also be a faster algorithm to recognize whether a given set of vertices K of a graph G is a potential maximal clique of G than the one of Corollary 12. We feel that improving these two points might lead to a much better time complexity for the treewidth and minimum fill-in algorithm.

A class of graphs may have an exponential number of minimal separators and consequently an exponential number of potential maximal cliques. Notice that there is no such class of graphs for which the treewidth problem has been solved in polynomial time, except the graphs of bounded treewidth. For example, the problem is still open for planar graphs. We think that a polynomial number of well-chosen potential maximal cliques could permit to compute or at least approximate the treewidth for classes of graphs with “many” minimal separators.

Acknowledgements

Thank you for the program of the Academic Leadership Grant (ALG), Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran (Indonesia), which has been providing facilities to conduct a research and publication.

References

- S. Arnborg, D.G. Corneil, A. Proskurowski, Complexity of finding embeddings in a k -tree, *SIAM J. Algebraic Discrete Methods* 8 (1987) 277–284.
- S. Arnborg, B. Courcelle, A. Proskurowski, D. Seese, An algebraic theory of graph reduction, *J. ACM* 40(1993) 1134–1164.
- S. Arnborg, A. Proskurowski, Linear time algorithms for NP-hard problems restricted to partial k -trees, *Discrete Appl. Math.* 23 (1989) 11–24.
- A. Berry, J.P. Bordat, O. Cogis, in: Generating all the minimal separators of a graph, *Workshop on Graph-theoretic Concepts in Computer Science (WG'99), Lecture Notes in Computer Science, Vol. 1665*, Springer, Berlin, 1999.
- H. Bodlaender, A tourist guide through treewidth, *Acta Cybernet.* 11 (1993) 1–23.
- H. Bodlaender, A linear-time algorithm for finding tree-decompositions of small treewidth, *SIAM J. Comput.* 25 (1996) 1305–1317.
- H. Bodlaender, Treewidth: Algorithmic techniques and results, *Proc. of MFCS'97, Lecture Notes in Computer Science, Vol. 1295*, Springer, Berlin, 1997, pp. 19–36.
- H. Bodlaender, B. de Fluiter, Reduction algorithms for constructing solutions of graphs with small treewidth. *Proc. COCOON'96, Lecture Notes in Computer Science, Vol. 1090*, Springer, Berlin, 1996, pp. 199–208.
- H. Bodlaender, J.R. Gilbert, H. Hafsteinsson, T. Kloks, Approximating treewidth, pathwidth, and minimum elimination tree height, *J. Algorithms* 18 (1995) 238–255.

- H. Bodlaender, J. Gustedt, J.A. Telle, Linear-time register allocation for a fixed number of registers. *Proc. 9th Annu. ACM-SIAM Symp. on Discrete Algorithms, San Francisco, CA, 1998*, ACM, New York, 1998, pp. 574–583.
- V. Bouchitté, I. Todinca, Minimal triangulations for graphs with “few” minimal separators, *Proc. 6th Annu. European Symp. on Algorithms (ESA'98), Lecture Notes in Computer Science, Vol. 1461*, Springer, Berlin, 1998, pp. 344–355.
- V. Bouchitté, I. Todinca, Treewidth and minimum fill-in of weakly triangulated graphs, *Proc. 16th Symp. on Theoretical Aspects of Computer Science (STACS'99), Lecture Notes in Computer Science, Vol. 1563*, Springer, Berlin, 1999, pp. 197–206.
- M.S. Chang, Algorithms for maximum matching and minimum fill-in on chordal bipartite graphs, *ISAAC'96, Lecture Notes in Computer Science, Vol. 1178*, Springer, Berlin, 1996, pp. 146–155.
- B. Courcelle, The monadic second-order logic of graphs III: treewidth, forbidden minors and complexity issues, *Informatique Théorique* 26 (1992) 257–286
- B. Courcelle, M. Mosbah, Monadic second-order evaluations on tree-decomposable graphs, *Theoret. Comput. Sci.* 109 (1993) 49–82.
- M.C. Golumbic, Algorithmic Graph Theory and Perfect Graphs, *Academic Press*, New York, 1980.
- T. Hagerup, Dynamic algorithms for graphs of bounded treewidth, *Proc. 24th Internat. Colloq. On Automata, Languages, and Programming (ICALP'97), Lecture Notes in Computer Science*, Springer, Berlin, 1997, pp. 292–302.
- S. Kannan, T. Warnow, Inferring evolutionary history from DNA sequences, *Proc. 31st Annu. Symp. on Foundations of Computer Science*, 1990, pp. 362–371.
- T. Kloks, H.L. Bodlaender, H. Müller, D. Kratsch, Computing treewidth and minimum fill-in: all you need are the minimal separators. *Proc. 1st Ann. European Symp. on Algorithms (ESA'93), Lecture Notes in Computer Science, Vol. 726*, Springer, Berlin, 1993, pp. 260–271.
- T. Kloks, H.L. Bodlaender, H. Müller, D. Kratsch, Erratum to the ESA'93 proceedings, *Proc. 2nd Annu. European Symp. on Algorithms (ESA'94), Lecture Notes in Computer Science, Vol. 855*, Springer, Berlin, 1994, p. 508.
- T. Kloks, D. Kratsch, Treewidth of chordal bipartite graphs, *J. Algorithms* 19 (2) (1995) 266–281.
- T. Kloks, D. Kratsch, Listing all minimal separators of a graph, *SIAM J. Comput.* 27 (3) (1998) 605–613.
- T. Kloks, D. Kratsch, H. Müller, Approximating the bandwidth for asteroidal triple-free graphs, *Proc. 3rd Annu. European Symp. on Algorithms (ESA'95), Lecture Notes in Computer Science, Vol. 979*, Springer, Berlin, 1995, pp. 434–447.
- T. Kloks, D. Kratsch, J. Spinrad, On treewidth and minimum fill-in of asteroidal triple. *Theoret. Comput. Sci.* 175 (1997) 309–335.
- T. Kloks, D. Kratsch, C.K. Wong, Minimum fill-in of circle and circular-arc graphs, *J. Algorithms* 28 (2) (1998) 272–289.
- J.W.H. Liu, The multifrontal method for sparse matrix solution: theory and practice, *SIAM Rev.* 34 (1) (1992) 82–109.
- A. Parra, P. ScheLer, Characterizations and algorithmic applications of chordal graph embeddings, *Discrete Appl. Math.* 79 (1–3) (1997) 171–188.
- N. Robertson, P. Seymour, Graphs minors. III. Planar tree-width, *J. Combin. Theory Ser. B* 36 (1984) 49–64.
- N. Robertson, P. Seymour, Graphs minors. II. Algorithmic aspects of tree-width, *J. Algorithms* 7 (1986) 309–322.
- D.J. Rose, Triangulating graphs and the elimination process, *J. Math. Anal. Appl.* 32 (1970) 597–609.
- R. Sundaram, K. Sher Singh, C. Pandu Rangan, Treewidth of circular-arc graphs, *SIAM J. Discrete Math.* 7 (1994) 647–655.
- M. Thorup, All structured programs have small tree width and good register allocation, *Inform. And Comput.* 142 (2) (1998) 159–181.
- I. Todinca, Aspects algorithmiques des triangulations minimales des graphes, *Ph.D. Thesis, École Normale Supérieure de Lyon*, 1999.
- M. Yannakakis, Computing the minimum fill-in is NP-complete, *SIAM J. Algebraic Discrete Methods* 2 (1981) 77–79.

Biographies

Mochamad Suyudi, is a lecturer at the Department of Mathematics, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran. Bachelor in Mathematics at the Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, and Master in Mathematics at the Faculty of Mathematics and Natural

Sciences, Universitas Gajah Mada. Currently pursuing Ph.D. program in the field of Graphs at Universiti Sultan Zainal Abidin (UNISZA) Malaysia Terengganu.

Sukono is a lecturer in the Department of Mathematics, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran. Currently serves as Head of Master's Program in Mathematics, the field of applied mathematics, with a field of concentration of financial mathematics and actuarial sciences.

Abdul Talib Bon is a professor of Production and Operations Management in the Faculty of Technology Management and Business at the Universiti Tun Hussein Onn Malaysia since 1999. He has a PhD in Computer Science, which he obtained from the Universite de La Rochelle, France in the year 2008. His doctoral thesis was on topic Process Quality Improvement on Beltline Moulding Manufacturing. He studied Business Administration in the Universiti Kebangsaan Malaysia for which he was awarded the MBA in the year 1998. He's bachelor degree and diploma in Mechanical Engineering which he obtained from the Universiti Teknologi Malaysia. He received his postgraduate certificate in Mechatronics and Robotics from Carlisle, United Kingdom in 1997. He had published more 150 International Proceedings and International Journals and 8 books. He is a member of MSORSM, IIF, IEOM, IIE, INFORMS, TAM and MIM.

Mustafa Mamat is a lecturer in the Faculty of Informatics and Computing, Universiti Sultan Zainal Abidin, Malaysia. Currently serves as Dean of Graduate School Universiti Sultan Zainal Abidin, Terengganu, Malaysia. The field of applied mathematics, with a field of concentration of optimization.